

Polysiloxane ionic liquids as good solvents for β-cyclodextrin-polydimethylsiloxane polyrotaxane structures

Narcisa Marangoci¹, Rodinel Ardeleanu¹, Laura Ursu¹, Constanta Ibanescu^{1,2}, Maricel Danu¹, Mariana Pinteala^{*1} and Bogdan C. Simionescu^{1,2}

Full Research Paper	Open Access
Address: ¹ Centre of Advanced Research in Bionanoconjugates and Biopolymers, "Petru Poni" Institute of Macromolecular Chemistry,	<i>Beilstein J. Org. Chem.</i> 2012 , <i>8</i> , 1610–1618. doi:10.3762/bjoc.8.184
700487 Iasi, Romania and ² Department of Natural and Synthetic	Received: 31 May 2012
Polymers, "Gheorghe Asachi" Technical University of Iasi, 700050	Accepted: 17 August 2012
lasi, Romania	Published: 24 September 2012
Email: Mariana Pinteala [*] - ninteala@icmon ro	This article is part of the Thematic Series "Superstructures with cyclodextrins: Chemistry and Applications"
Manana i inteala - pinteala@ichipp.ro	
* Corresponding author	Guest Editor: H. Ritter
Keywords: cyclodextrins; imidazolium salt; ionic liquid; polyrotaxanes; polysiloxanes	© 2012 Marangoci et al; licensee Beilstein-Institut. License and terms: see end of document.

Abstract

An ionic liquid based on polydimethylsiloxane with imidazolium salt brushes was synthesized as a good solvent for β -cyclodextrinpolydimethylsiloxane rotaxane. As expected the PDMS-Im/Br ionic liquid had a liquid-like non-Newtonian behavior with rheological parameters dependent on frequency and temperature. The addition of rotaxane to the ionic liquid strengthened the non-Newtonian character of the sample and a type of stable liquid-like network was formed due to the contribution of weak ionic interactions. The structure is stable in the 20 to 80 °C domain as proved by the oscillatory and rotational rheological tests.

Introduction

Ionic liquids (ILs) are environmentally friendly solvents with great potential for chemical and nonchemical applications due to their low melting points, nonvolatile and noncorrosive properties at room temperature. They possess good conductivities [1,2] and often they present a good thermal stability up to 400 $^{\circ}$ C [1,3]. The potential applications of ILs made them the

subject of a number of works that showed their use as solvents or as solvents for synthesis and catalysis [4-6]. In this context ILs, which have polar and nonpolar regions, could play an important role in the field of supramolecular organization of different supramolecular structures (such as polyrotaxanes or supermolecules formed by ILs with different host molecules),